

*Hadronic Contributions to R and $g - 2$ from
Initial-State-Radiation Data*



Denis Bernard

Laboratoire Leprince-Ringuet, Ecole polytechnique IN2P3/CNRS

Palaiseau, France

On behalf of the *BABAR* Collaboration


Physics in Collision - 31 Aug 2009 - Kobe, Japan

The “Anomalous” Magnetic Moment of the Lepton

$$\vec{\mu} = g \frac{e}{2m} \vec{s}, \quad a = (g - 2)/2$$

(1928) Pointlike Dirac particles : $g = 2$, $a = 0$.

$g \neq 2$ due to higher order contributions :



- (1947) Nafe et al. measure
- (1948) Schwinger (1st order)

$$a_e = (2.6 \pm 0.5) \times 10^{-3}$$


$$a^{(1)} = \alpha/2\pi \approx 1.2 \times 10^{-3}$$

Lepton universality at this 1st order

Our belief in QED and in the gauge-theory-based SM originates from this 1st success.


Higher Orders

One graph given as example out of many ..




$$a = a^{\text{QED}} + a^{\text{had}} + a^{\text{weak}}$$


Hadronic Vacuum Polarisation
(VP)



Hadronic light-by-light
Scattering



Weak
Interactions



a_e, α and a_μ

- Heavy-to-Light and Light-to-Heavy mass ratios take part differently (e/μ) in the loops (QED, QCD, weak)

$$a_e = \frac{\alpha}{2\pi} - 0.3 \left(\frac{\alpha}{\pi}\right)^2 + 1.2 \left(\frac{\alpha}{\pi}\right)^3 - 1.9 \left(\frac{\alpha}{\pi}\right)^4 + 0.0(4.6) \left(\frac{\alpha}{\pi}\right)^5 + 1.72(2)10^{-12}(\text{QCD} + \text{weak})$$
$$a_\mu = \frac{\alpha}{2\pi} + 0.8 \left(\frac{\alpha}{\pi}\right)^2 + 24. \left(\frac{\alpha}{\pi}\right)^3 - 131. \left(\frac{\alpha}{\pi}\right)^4 + 663. \left(\frac{\alpha}{\pi}\right)^5 + 7.07(7)10^{-8}(\text{QCD} + \text{weak})$$

Numbers truncated !

- a_e measured in a one-electron quantum cyclotron



$$a_e = 1159652180.73(\pm 0.28)10^{-12}, \quad (0.24 \text{ ppb})$$

- $\Rightarrow \alpha$ known to 0.37 ppb
- In total the QED uncertainty on a_μ^{QED} is tiny : 1.7 ppb

Odom PRL 97 (2006) 030801,

Gabrielse PRL 97 (2006) 030802

a_μ Measurement



- $\pi^- \rightarrow \mu^- \nu$ violates P, μ^- longitudinally polarized.
- μ^- stored in a cyclotron, constant \vec{B} .
 - μ^- rotating with freq ω_c ; μ^- spin precessing with freq ω_s
 - freq. difference $\omega_a = \omega_s - \omega_c = a_\mu eB/m_\mu$
- $\mu \rightarrow e \nu \bar{\nu}$ violates P, e direction (energy in lab) remembers μ^- polarization.
- Fraction of e above $E_{\text{threshold}}$ is modulated with freq. ω_a

$$a_\mu(\text{expt}) = (11659208.0 \pm 5.4(\text{stat}) \pm 3.3(\text{syst})) \times 10^{-10}, \quad (0.54 \text{ ppm})$$

Theoretical prediction for a_μ – May 2009

SM-to-experiment comparison [in units 10^{-10}]


QED	116 584 71.81	± 0.02
Leading hadronic vacuum polarization (VP)	690.30	± 5.26
Sub-leading hadronic vacuum polarization	-10.03	± 0.11
Hadronic light-by-light	11.60	± 3.90
Weak (incl. 2-loops)	15.32	± 0.18
<hr/>		
Theory	11659179.00	± 6.46
Experiment	11659208.00	± 6.30
<hr/>		
Exp – theory	29.00	± 9.03

Assuming Gaussian statistics, a 3.2σ discrepancy.

Jegerlehner, Nyffeler / Phys Rept 477 (2009) 1110

uses e^+e^- input only for VP

Theoretical prediction : The Hadronic VP (1)



- Quark loops not computable from first principles – QCD.
- Vacuum polarization : energy dependent running charge :

$$e^2 \rightarrow e^2 / [1 + (\Pi'(k^2) - \Pi'(0))]$$


- Dispersion relation from analyticity

$$\Pi'(k^2) - \Pi'(0) = \frac{k^2}{\pi} \int_0^\infty \frac{Im\Pi'(s)}{s(s - k^2 - i\epsilon)} ds$$

- Optical theorem (unitarity)

$$Im\Pi'(s) = \alpha(s)R_{\text{had}}(s)/3, \quad \text{with } R_{\text{had}}(s) = \sigma_{\text{had}} \frac{3s}{4\pi\alpha(s)} = \frac{\sigma_{e^+e^- \rightarrow \text{hadrons}}}{\sigma_{e^+e^- \rightarrow \mu^+\mu^-}}$$

Theoretical prediction : The Hadronic VP (2)




Wrapping it up, the “dispersion integral” :

$$a_\mu^{\text{had}} = \left(\frac{\alpha m_\mu}{3\pi} \right)^2 \int \frac{R_{\text{had}}(s) \hat{K}(s)}{s^2} ds$$

- Technically, $\int_{4m_\pi^2}^{E_{\text{cut}}^2}$ is obtained from the data, $\int_{E_{\text{cut}}^2}^{\infty}$ from pQCD.
- The estimation of the contribution with the largest uncertainty to $a_\mu(\text{theory})$ boils down to a precise measurement of $R_{\text{had}}(s)$
- Most precision on $R_{\text{had}}(s)$ needed at low \sqrt{s}


$R_{\text{had}}(s)$: Direct Measurements $e^+e^- \rightarrow \text{Hadrons}$



PDG, Phys. Lett. B667,1 (2008)

$e^+e^- \rightarrow \text{Hadrons} : \text{channel break-down}$

Contributions to the “dispersion integral”.




$e^+e^- \rightarrow \pi^+\pi^-$ dominates (73 %)

$$a_{\mu}^{\pi^+\pi^-} [2m_{\pi}, 1.8 \text{ GeV}/c^2] = (504.6 \pm 3.1(\text{exp}) \pm 0.9(\text{rad})) \times 10^{-10}$$

$$a_{\mu}^{\text{had}} = (690.9 \pm 5.3) \times 10^{-10}$$

Davier, Nucl. Phys. Proc. Suppl. 169, 288 (2007)

$e^+e^- \rightarrow \pi^+\pi^-$: Direct Measurements




- (KLOE 08 supersedes KLOE04)
- The 3.2σ discrepancy mentioned above is based on this input

Jegerlehner, Nyffeler / Phys Rept 477 (2009) 1110



τ Decay Spectral Functions

$I = 1$ part of $e^+e^- \rightarrow$ had from $\tau \rightarrow \nu_\tau +$ had by isospin rotation



- example $\pi^0\pi^- \leftrightarrow \pi^+\pi^-$
- $\sigma(e^+e^- \rightarrow \pi^+\pi^-) = \frac{4\pi\alpha^2}{s}\nu_0(s),$ $\nu(s)$ “spectral function”
- $\frac{1}{\Gamma} \frac{d\Gamma}{ds} = F(s) \frac{\mathcal{B}(\tau \rightarrow e\nu_\tau\bar{\nu}_e)}{\mathcal{B}(\tau \rightarrow \pi^-\pi^0\nu_\tau)} \times \nu_-(s),$ where $F(s)$ is a known function of s
- CVC : $\nu_0(s) = \nu_-(s)$ isospin breaking (IB) corrections ...
- ALEPH (1997), OPAL (1999), CLEO (2000)
- $\mathcal{B}(\tau \rightarrow \pi^-\pi^0\nu_\tau)$ ALEPH’s most precise $(25.471 \pm 0.097 \pm 0.085)\%$

Belle's High Statistics Results on $\tau \rightarrow \pi^- \pi^0 \nu_\tau$



	$a_{\mu}^{\pi^+\pi^-} [2m_{\pi}, 1.8 \text{ GeV}/c^2] \text{ (units } 10^{-10})$
ALEPH, CLEO, OPAL	$520.1 \pm 2.4 \text{ (exp)} \pm 2.7 \text{ (Br.)} \pm 2.5 \text{ (IB)}$
Belle	$523.5 \pm 1.5 \text{ (exp)} \pm 2.6 \text{ (Br.)} \pm 2.5 \text{ (IB)}$


72 fb^{-1} of e^+e^- data taken at $\approx 10.6 \text{ GeV}$



Fujikawa Phys.Rev.D78 :072006,2008.

Combination from Davier Eur. Phys. J. C 27, 497 (2003).

τ Spectral Functions : Isospin Breaking (IB) Corrections



Several contributions to the IB corrections, among which :

- Short- and long-distance radiative corrections
- $\pi^0 - \pi^\pm$ and $\rho^0 - \rho^+$ mass differences
- FSR correction included


IB corrections are still being worked on :

$$-13.8 \pm 2.4 + 4.2(\text{FSR}) \quad \text{Eur. Phys. J. C 27, 497 (2003).}$$

$$-16.1 \pm 1.5, \quad \text{0906.5443v2 [hep-ph] sub. to Eur. Phys. J. C}$$

$$\Delta = -6.9 \quad \text{Units } 10^{-10}$$

Initial State Radiation (ISR)



- Optimal use of the available luminosity
- Covers whole energy range with same detector condition and analysis.
 - And good efficiency down to threshold
- If observe the whole final state ($\gamma + \text{hadrons}$)
 \Rightarrow over-constrained kinematical fit \Rightarrow powerful background noise rejection.

$$\frac{d\sigma_{[e^+e^- \rightarrow f\gamma]}(s')}{ds'} = \frac{2m}{s} W(s, x) \sigma_{[e^+e^- \rightarrow f]}(s') , \quad x = \frac{E_\gamma}{\sqrt{s}} = 1 - \frac{s'}{s}$$

- $W(s, x)$ “radiator function”, density of probability to radiate a photon with energy $E_\gamma = x\sqrt{s}$: a known function [Binner, Physics Letters B 459 \(1999\) 279](#)

ISR : “Old” BaBar Results

Vigourous campaign that is still in progress

$K^+K^-\eta, K^+K^-\pi^0, K^0K^\pm\pi^\mp$

Phys.Rev.D77 :092002,2008.

$2(\pi^+\pi^-)\pi^0, 2(\pi^+\pi^-)\eta, K^+K^-\pi^+\pi^-\pi^0, K^+K^-\pi^+\pi^-\eta$

Phys.Rev.D76 :092005,2007.

$K^+K^-\pi^+\pi^-, K^+K^-\pi^0\pi^0, K^+K^-\pi^+\pi^-\pi^0$

Phys.Rev.D76 :012008,2007.

$3(\pi^+\pi^-), 2(\pi^+\pi^-\pi^0), K^+K^-\pi^+\pi^-\pi^0$

Phys.Rev.D73 :052003,2006.

$\bar{p}p$

Phys.Rev.D73 :012005,2006.

$2(\pi^+\pi^-), K^+K^-\pi^+\pi^-, K^+K^-\pi^+\pi^-\pi^0$

Phys.Rev.D71 :052001,2005.

$\pi^+\pi^-\pi^0$

Phys.Rev.D70 :072004,2004.

- First observations

232 fb⁻¹, 89 fb⁻¹ @ 10.6 GeV

- ISR γ tagging \Rightarrow efficient background rejection
- Only charmless mesons in this slide
- Unprecedented accuracy :




		without <i>BABAR</i>	with <i>BABAR</i>
$a_\mu(< 1.8 \text{ GeV}/c^2)$	$\pi^+\pi^-\pi^0$	2.45 ± 0.26	3.25 ± 0.09
	$2(\pi^+\pi^-)$	14.20 ± 0.90	13.09 ± 0.44
	$3(\pi^+\pi^-)$	0.10 ± 0.10	0.11 ± 0.02
	$2(\pi^+\pi^-\pi^0)$	1.42 ± 0.30	0.89 ± 0.09

Davier, Nucl. Phys. Proc. Suppl. 169, 288 (2007)

ISR : KLOE $\pi^+ \pi^-$

- 240 pb^{-1}
- On the ϕ
- No ISR photon tagging
- \vec{p}_{had} compatible with a γ in beam pipe
- LO : No additional photon reco'ed
- $W(s, x)$ from NLO PHOKHARA generator (precision 0.5 %)
- Luminosity (0.3 %) from Bhabha scat. (BABAYAGA generator (0.1 %))



$$a_{\mu}^{\pi^+\pi^-} [592 - 975 \text{ MeV}] = (387.2 \pm 0.5(\text{stat}) \pm 2.4(\text{expt}) \pm 2.3(\text{th})) \times 10^{-10}$$

- Compatible with CMD-2 & SND on (630 - 958 MeV) with similar precision : $\Delta = (-4.6 \pm 4.2) \times 10^{-10}$



Ambrosino, Phys.Lett.B670 :285,2009

ISR : BaBar $\pi^+ \pi^-$

Only attempt in BaBar, to my knowledge, to master systematics at the 10^{-3} level

- ISR γ in EMC (thus : at large angle)
- 1 (for efficiency) or 2 (for physics) tracks of good quality
- Particle identification (PID) of the charged particles
- separate $\pi\pi$, KK , $\mu\mu$ event samples
- kinematic fit (using only direction of ISR γ) including 1 additional γ : NLO !
- obtain all efficiencies (trigger, filter, tracking, PId, fit) from same data
- measure ratio of $\pi\pi$ to $\mu\mu$ cross sections to cancel : ee luminosity, additional ISR, vacuum polarization, ISR γ efficiency

Correct for FSR in $\mu\mu$ and ISR + additional FSR, both calc. in QED, and checked in data

$$R_{\text{exp}}(s') = \frac{\sigma_{[\pi\pi\gamma(\gamma)]}(s')}{\sigma_{[\mu\mu\gamma(\gamma)]}(s')} = \frac{\sigma_{[\pi\pi(\gamma)]}^0(s')}{(1 + \delta_{\text{FSR}}^{\mu\mu})\sigma_{[\mu\mu(\gamma)]}^0(s')} = \frac{R(s')}{(1 + \delta_{\text{FSR}}^{\mu\mu})(1 + \delta_{\text{add,FSR}}^{\mu\mu})}$$

A Comment : γ -Tag or not γ -Tag ?



A key issue is the difficulty (impossibility?) to control the ISR γ efficiency to the desired precision.

Two ways out :

- KLOE : no γ tag, at the cost of a significant background.
(mitigated by requesting that the non-observed γ would be in the beam pipe)
- BaBar : γ tag, and use the $\pi\pi/\mu\mu$ ratio
 - \Rightarrow the ISR γ efficiency cancels in the ratio (to first order)
 - \Rightarrow γ tag costs a loss of 9/10 in statistics

Note that KLOE has a tagged analysis in progress.

ISR : BaBar $\pi^+ \pi^-$



Bare (VP removed), unfolded $\sigma_{e^+e^- \rightarrow \pi^+\pi^-}$ 232 fb^{-1} @ $\sqrt{s} \approx 10.6 \text{ GeV}$



$$a_{\mu}^{\pi^+\pi^-} [2m_{\pi}, 1.8 \text{ GeV}/c^2] = (514.1 \pm 2.2 \pm 3.1) \times 10^{-10}$$


- Similar precision as combination of previous e^+e^- results.
- 2.0 σ larger than previous e^+e^- average submitted to PRL [arXiv :0908.3589v1 \[hep-ex\]](https://arxiv.org/abs/0908.3589v1)
- Longer paper, in preparation, [to be submitted to PRD](#)

Systematics

Relative systematic uncertainties (in 10^{-3}) on the $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ cross section by $\sqrt{s'}$ intervals (in GeV) up to 1.2 GeV.

Source of Uncertainty	CM Energy Interval (GeV)				
	0.3-0.4	0.4-0.5	0.5-0.6	0.6-0.9	0.9-1.2
trigger/ filter	5.3	2.7	1.9	1.0	0.5
tracking	3.8	2.1	2.1	1.1	1.7
π -ID	10.1	2.5	6.2	2.4	4.2
background	3.5	4.3	5.2	1.0	3.0
acceptance	1.6	1.6	1.0	1.0	1.6
kinematic fit (χ^2)	0.9	0.9	0.3	0.3	0.9
correlated $\mu\mu$ ID loss	3.0	2.0	3.0	1.3	2.0
$\pi\pi/\mu\mu$ non-cancel.	2.7	1.4	1.6	1.1	1.3
unfolding	1.0	2.7	2.7	1.0	1.3
ISR luminosity ($\mu\mu$)	3.4	3.4	3.4	3.4	3.4
total uncertainty	13.8	8.1	10.2	5.0	6.5


BaBar : Sanity check : Comparison of the $\mu\mu$ spectrum with QED



- Here the radiator function is needed.
- MC simulation corrected for all known MC/data differences.
 - e.g. : ISR γ efficiency measured in data, from $\mu\mu$ -only reco'ed evts.
 - MC corrected for known NLO deficiencies by comparing to PHOKHARA

Good agreement within $0.4 \pm 1.1\%$; dominated by $\mathcal{L}_{e^+e^-}$ ($\pm 0.9\%$)

VDM Fit of $|F_\pi(s)|^2$




- $|form\ factor|^2$ fitted with a vector dominance model, $\rho, \rho', \rho'', \omega$.
- ρ 's described by the Gounaris-Sakurai model $\chi^2/n_{df} = 334/323$




BaBar $\pi^+\pi^-$: comparison with previous results


Belle τ




CMD2 e^+e^-



KLOE e^+e^- ISR




SND e^+e^-



The green band is the representation of the VDM fit to the BaBar ISR data

a_μ : Where do we stand ?



- Good agreement between e^+e^- -based predictions
- BNL a_μ measurement 3.4σ higher than e^+e^- -based
- τ -based “only” 1.8σ from BNL.

My combinations, assuming Gaussian statistics : back-up slides

- BaBar compatible with the τ -based
- BaBar 2.4σ from BNL.
- $(e^+e^- + \text{BaBar})$ 3.3σ from BNL.

The $(e^+e^- - \text{BNL})$ shift has decreased, so has the uncertainty .. $> 3\sigma$ still

What could happen during the next decade

- $a_\mu(\text{expt}) : 0.54 \rightarrow 0.14 \text{ ppm}$ Carey, FERMILAB-PROPOSAL-0989 (2009)
 - was statistics limited (0.46 ppm); move μ storage ring BNL \rightarrow FNAL.
- Vacuum polarization (hadronic) : $a_\mu^{\pi^+\pi^-}$ is and will remain the bottle-neck
 - e^+e^- , “direct”
 - upgrades, CMD2 \rightarrow CMD3 and SDN
 - BES III : $R : 2.0 - 4.6 \text{ GeV}$, Zhemchugov @ TAU08
 - e^+e^- , ISR : Statistics is not an issue.
 - Belle checks BaBar’s ISR result ?
 - KLOE checks it’s ISR result with $\pi\pi/\mu\mu$ ratio
 - τ : Statistics is not an issue.
 - Theorists converge to a narrow range calculation of ΔIB
 - BES III : τ spectral fns, $\mathcal{B}(\tau \rightarrow \pi^-\pi^0\nu)$ Zhemchugov @ TAU08
 - BaBar checks Belle’s τ result ?
- Light-by-Light : Theory + $\gamma\gamma$ program at DAΦNE-2 Prades @ TAU08
- QED : The 5th order is being evaluated Kinoshita 先生

ご清聴ありがとうございました。

Je vous remercie de votre attention distinguée

Back-up slides

Recent Improvements

		Factor	
Harvard Measurement of a_e (2006)	Odom	15	(1987) Van Dyck
BNL Measurement of a_μ (2006)	Bennett	14	(1976) Bailey

Jargon

ISR Initial State Radiation

FSR Final State Radiation

VP Vacuum Polarization

IB Isospin Breaking

CVC Conserved vector current

P Parity

pQCD perturbative QCD

VDM Vector dominance model

My combinations

- I assume Gaussian statistics
- I use the combination of former e^+e^- results with the recent ISR KLOE result, as performed by Davier09, i.e. (All in units 10^{-10})
- Care is taken of correlations in combinations.

$$a_{\mu}^{\pi^+\pi^-} [2m_{\pi}, 1.8 \text{ GeV}/c^2] = 503.5 \pm 3.5$$

$$a_{\mu} = 11659177.7 \pm 4.4 \pm 2.6$$

- The combination of the former $a_{\mu}^{\pi^+\pi^-} [2m_{\pi}, 1.8 \text{ GeV}/c^2]$ and of BaBar's result is 508.3 ± 2.6 .
- Replacing the former e^+e^- results by the Babar one, $11\ 659\ 188.3 \pm 5.3$
- Combining the former e^+e^- results and the Babar one, $11\ 659\ 182.5 \pm 4.5$

Slide added after my talk

- Concerning the minor differences between the numerical values of combined results I gave, with respect with those of Boris Shwartz' talk at LP09 :
Boris was refering to version 1 of arXiv :0906.5443 [hep-ph] while I am refering to version 2.
- After my talk, Changzheng Yuan has advertized a soon-to-appear preprint on "Reevaluation of the hadronic contribution to the muon magnetic anomaly using new $e^+e^- \rightarrow \pi^+\pi^-$ cross section data from BABAR"
Here it is : arXiv :0908.4300 [hep-ph]
- After my talk, Naohito Saito has mentioned a proposal for an alternative (to the BNL scheme) design for a new $(g - 2)$ experiment
See slides 46 – 53 of his talk on Spin Physics.