



# Color screening in Quark Gluon Plasma (QGP): A new experiment to measure charm production in PbPb collisions at the CERN SPS

CHIC: **C**harm in **H**eavy **I**on **C**ollisions

*F. Fleuret <sup>1</sup>, F. Arleo <sup>2</sup>, E. G. Ferreiro <sup>3</sup>, P.-B. Gossiaux <sup>4</sup>, S. Peigné <sup>4</sup>*

<sup>1</sup> LLR, École polytechnique – IN2P3/CNRS, Palaiseau, France

<sup>2</sup> LAPTh, Université de Savoie – CNRS, Annecy-le-Vieux, France

<sup>3</sup> Universidad de Santiago de Compostela, Santiago de Compostela, Spain

<sup>4</sup> SUBATECH, université de Nantes – IN2P3/CNRS, Nantes, France


## • Heavy quarks and Quark Gluon Plasma

**Heavy quarks are "special" QGP probes** :  $m_Q \gg$  QGP critical temperature  $T_c$  ( $\sim 170$  MeV),

- Heavy quarks should be produced in initial hard nucleon-nucleon collisions only, the QGP phase shouldn't modify the overall heavy quark yields,
- On the other hand, The **QGP phase should modify  $Q\bar{Q}$  bound state yields**

### – Possible QGP effects:

- **Color screening:  $Q\bar{Q}$  bound states suppression**
  - Color screening in a QGP decreases quarkonium binding
  - Color screening should lead to a suppression of quarkonium production yields
- **Recombination:  $Q\bar{Q}$  bound states enhancement**
  - at sufficiently high energies,  $Q\bar{Q}$  pairs are abundantly produced.
  - Statistical combination can lead to an enhancement of quarkonium production yields



- Experimentally, charmonium is a privileged probe

- Charmonium production in A+A collisions studied at:


- |            |                        |                              |
|------------|------------------------|------------------------------|
| • CERN-SPS | ( $\sqrt{s}=17$ GeV)   | NA38, NA50, NA60 experiments |
| • BNL-RHIC | ( $\sqrt{s}=200$ GeV)  | PHENIX, STAR experiments     |
| • CERN-LHC | ( $\sqrt{s}=2.76$ TeV) | ALICE, CMS experiments       |

- Short summary for  $J/\Psi$ :

- |                      |   |
|----------------------|---|
| • NA50 (PbPb@SPS)    | observed an <i>anomalous</i> $J/\Psi$ suppression |
| • PHENIX (AuAu@RHIC) | observed a similar suppression (than NA50)        |
| • ALICE (PbPb@LHC)   | observed a smaller suppression (than PHENIX)      |

- ➔ Possible Color screening starting at SPS
- ➔ Possible recombination occurring at LHC

- Within the SPS+RHIC+LHC energy range, charm seems to be the adequate probe to investigate both screening and recombination.



## • What next to be done with charmonium

To confirm (and study) charmonium color screening and enhancement, one must compare charmonium and open charm production in A+A collisions

- Since most of the produced  $c\bar{c}$  pairs hadronize into open charm, open charm production reflects the original  $c\bar{c}$  pair production
- Open charm is therefore an (the?) appropriate reference to calibrate charmonium screening/recombination studies.

### – Study charmonium recombination


- Both  $J/\Psi$  and open charm will be measured in PbPb at large energy densities at LHC
- **LHC is the best place to study recombination**

### – Study charmonium color screening

- At SPS energies, in Pb+Pb collisions,  $J/\Psi$  suppression occurs in the middle of the accessible energy density range

→ **SPS is the best place to study color screening**

- *Need measurement of open charm yields*
- *Need precise measurements of several  $c\bar{c}$  states to test if color screening leads indeed to a sequential suppression*



- **Quarkonium sequential suppression**

- **Quarkonium sequential suppression** in a Quark Gluon Plasma is a prediction of **lattice QCD**, for instance :

[H. Satz, J. Phys. G 32 \(2006\)](#)

quarkonium dissociation temperature  
critical QGP temperature →

state	$J/\psi(1S)$	$\chi_c(1P)$	$\psi'(2S)$	$\Upsilon(1S)$	$\chi_b(1P)$	$\Upsilon(2S)$	$\chi_b(2P)$	$\Upsilon(3S)$
$T_d/T_c$	2.10	1.16	1.12	> 4.0	1.76	1.60	1.19	1.17

- Because of feed-downs and different  $T_d$ , **sequential suppression** should show up.

*Feed-downs contributing to  $J/\Psi$  inclusive yield*


**60% direct  $J/\Psi$**   
**+ 30%  $\chi_c \rightarrow J/\Psi + \gamma$**   
**+ 10%  $\Psi' \rightarrow J/\Psi + X$**

---

**Inclusive  $J/\Psi$  yield**

According to lattice calculations,  
 $T_d(\Psi') < T_d(\chi_c) < T_d(J/\Psi)$

→ **One should observe a step-like suppression pattern**



- Anomalous suppression at SPS**

[Eur.Phys.J.C49:559-567,2007](#)

$L$  = length of nuclear matter seen by quarkonium state

**Expected** = measured yields in p+A extrapolated to large  $L$


**Color screening ?**

NA50 measured  $J/\Psi$  and  $\Psi'$ , but,

- too small  $\Psi' \rightarrow J/\Psi$  feed-down
- too fragile  $\Psi'$

to answer the question

- $\rightarrow$  need of a larger feed-down fraction
- $\rightarrow$  Need of a stronger bound state
- $\rightarrow$  Need to measure  $\chi_c$  yield !



- Anomalous suppression at SPS

[Eur.Phys.J.C49:559-567,2007](http://arxiv.org/abs/hep-ex/0608007)

## Color screening ?


Take advantage of large  $\chi_c \rightarrow J/\Psi$  feed-down fraction

60% direct  $J/\Psi$   
 + 30%  $\chi_c \rightarrow J/\Psi + \gamma$   
 + 10%  $\Psi' \rightarrow J/\Psi + X$   
**Inclusive  $J/\Psi$  yield**

Measuring  $J/\Psi$ ,  $\Psi'$  and  $\chi_c$  suppression patterns will give the answer

- Alternative (no QGP) scenario: suppression by comoving hadrons

- Smooth suppression
- Same suppression-starting point
- Slopes related to binding energy :  $S_{\Psi'} > S_{\chi} > S_{J/\Psi}$



• **Must measure :**

- Charmonia :  $J/\Psi$ ,  $\Psi'$ ,  $\chi_c$
- open charm (for reference)

• **Beam:fixed-target experiment**

- high-intensity 158 GeV/c Pb beam
- high-intensity 158/450 GeV/c p beam

• **Experimental constraints**

- Measure muons from charmonia and open charm decays
- Measure photon from  $\chi_c$  decay ( $\chi_c \rightarrow J/\Psi + \gamma$ )

• **Detector main components :**

1. Vertex detector + Spectrometer


- Measures tracks before absorber  $\rightarrow$  very good mass resolution
- Measure muon vertex offset  $\rightarrow$  open charm

2. Ultra-granular calorimeter


- Measure  $\gamma$  in high  $\pi^0$  multiplicity environment

3. Absorber/ muon trigger

- Absorb  $\pi/K$
- Minimize fake triggers from  $\pi/K$  decays







**Calorimeter:**

- ultra-granular EMCal
- W + Si layers à la CALICE
  - 30 layers
  - 0.5 x 0.5 cm<sup>2</sup> pads
  - 24 X<sub>0</sub> in 20 cm
  - $\Delta E/E \sim 15\% / \sqrt{E}$



**Magnet :**  
1m long 2.5 T dipole

**Instrumented Absorber :**  
4.5 m thick Fe absorber  
→ dimuon trigger rate  $\sim 0.3$  kHz  
Could be magnetized to measure muon momentum

**Silicon Spectrometer**  
covers 1.5 rapidity unit  
 $\Delta p/p = 1\% \rightarrow J/\Psi$  mass resolution  $\sim 20$  MeV/c<sup>2</sup>

Estimations based on NA60/CERN telescope performances

- **Typical mass plots** (~1 week data taking w/ a 10%  $\lambda_1$  Pb target)
  - 200 000  $J/\Psi$  embedded in Pb+Pb Minbias events produced w/ EPOS
    - 140 000 direct  $J/\Psi \rightarrow \mu^+\mu^-$  (70%)
    - 60 000  $\chi_c \rightarrow J/\Psi \gamma \rightarrow \mu^+\mu^- \gamma$  (30%)



**After acceptance/selection cuts**  
 within  $y_{\text{CMS}} \in [-0.5; 0.5]$   
 35 000  $J/\Psi \rightarrow \mu^+\mu^-$   
**→ acc x eff = 17.4%**

*Including*  
 1700  $\chi_c \rightarrow J/\Psi \gamma \rightarrow \mu^+\mu^- \gamma$   
**→ acc x eff = 2.8 %**

- **Typical 40-day Pb+Pb run** ( $10^7 \cdot s^{-1}$  Pb beam  $\rightarrow$  10%  $\lambda_1$  Pb target)

- $\sim 180\,000$   $J/\Psi \rightarrow \mu^+\mu^-$  recorded

- 2 extreme *numerical* scenarios:

- If  $\chi_c$  suppressed as  $J/\Psi$   $\frac{\chi_c \text{ yield}}{J/\Psi \text{ yield}} \sim 4\%$


$$\rightarrow \left( \begin{array}{c} \text{most periph.} \\ \chi_c \text{ yield} \end{array} \right) = 16942 \times 4\% = 677$$

- If  $\chi_c$  suppressed as  $\Psi'$   $\frac{\chi_c \text{ yield}}{\Psi' \text{ yield}} = 2.18$

$$\left( \begin{array}{c} \text{most periph.} \\ \chi_c \text{ yield} \end{array} \right) = 16942 \times 4\% \times 0.6 = 406$$

$E_T$ range (GeV)	$\psi'$	$J/\psi$	$\chi_c$ as $\Psi'$	$\chi_c$ as $J/\Psi$
3–20	$186 \pm 25$	$16942 \pm 146$	406	677
20–35	$243 \pm 31$	$25229 \pm 181$	530	1010
35–50	$227 \pm 35$	$27276 \pm 192$	495	1091
50–65	$193 \pm 36$	$27681 \pm 196$	421	1107
65–80	$154 \pm 36$	$27315 \pm 200$	336	1093
80–95	$159 \pm 37$	$25111 \pm 193$	647	1004
95–150	$110 \pm 40$	$28570 \pm 209$	240	1143
			<b>3075</b>	<b>7125</b>



NA50 data



**Expect  $3000 < N_{\chi_c} < 7000$**

- Use same Strategy as NA60: measure muon vertex

– Open charm decay length:  $\begin{cases} D^{+/-}: c\tau = 311.8 \mu\text{m} \\ D^0: c\tau = 122.9 \mu\text{m} \end{cases}$



**CHIC: Vertex detector located 7.5 cm downstream of the target (7 cm for NA60)**




- **CHIC is able to measure open charm yields.**
- Detailed simulations needed to estimate performances

A thorough p+A program is mandatory to study Cold Nuclear Matter effects as a reference to study Hot Nuclear Matter effects


- **Must control (understand) :**

- charmonium absorption by cold nuclear matter → A dependence
  - Shadowing/anti-shadowing ( $x_2$  scaling)
  - Energy loss, formation time ( $x_F$  scaling)
- **Need large  $y_{\text{CMS}}$  range**


**Mid-rapidity :  $y_{\text{CMS}} \in [-0.5 ; 1]$**



**Forward-rapidity :  $y_{\text{CMS}} \in [0.5 ; 2]$**



- **Large rapidity range**
  - Significantly Larger rapidity range for CHIC ( $y_{CMS} \in [-0.5; 2]$ ).vs. NA50 ( $y_{CMS} \in [0; 1]$ )
- **Precise A dependence (thanks to fixed-target mode)**
  - NA50 samples : p+Be, p+Al, p+Cu, p+Ag, p+W, p+Pb
- **Large amount of data (thanks to fixed-target mode)**
  - **Large statistics** required to study  $J/\Psi$ ,  $\Psi'$ ,  $\chi_c$  and open charm differential yields as a function of  $y$ ,  $p_T$ .
  - **Current SPS operation: Delivering proton beam to the LHC several months per year**
  - **Significantly larger (than NA50) amount of data available for CHIC.**




Typical 1 week/target NA50 data taking (EPJ C33 (2004) 31-40)

Target	size ( $\lambda_I$ )	$\langle I_{protons} \rangle$ ( $\times 10^8$ )	Total $N_{protons}$ ( $\times 10^{12}$ )	$N_{\mu\mu}^{+-}$ (2.7 - 3.5)
Be	60 %	21.7	50.7	368 000
Al	52 %	23.0	63.4	602 000
Cu	28 %	27.0	45.5	762 000
Ag	30 %	24.8	43.8	821 000
W	19 %	23.5	28.5	524 000

# Expression of interest

- Submitted to SPSC (oct. 2012)

Expression of Interest  
Submitted to SPSC – oct.2012  
[CERN-SPSC-2012-031](#)



MINUTES of the 108th Meeting of the SPSC  
15-16 January 2013  
[CERN-SPSC-2013-008](#)



The SPSC has received an expression of interest to study charm production with proton and heavy ion beams. The SPSC recognizes the **strong physics motivation** of a study that addresses **central open questions** about the **color screening** of charmonium in heavy ion collisions and about **cold nuclear matter effects**. For a comprehensive investigation, an extension including open charm production would be desirable.

For further review, the SPSC would require a letter of intent with information about the experimental implementation and the **collaboration** pursuing it.

# Conclusion

- Measuring  $J/\Psi$ ,  $\Psi'$ ,  $\chi_c$  and open charm in A+A collisions at SPS will (dis)prove **sequential suppression scenario**.
  - Measuring  $J/\Psi$ ,  $\Psi'$ ,  $\chi_c$  and open charm in p+A collisions with several targets will give a thorough control of **Cold Nuclear Matter effects**
  - The apparatus is well suited to explore other important physics subjects such as low mass lepton pairs production in heavy ion collisions.
- Testing sequential suppression scenario at SPS is crucial to fully understand RHIC and LHC results.**



# Project status

- **Green light from CERN SPSC**
  - Eol submitted to SPSC in oct. 2012 : [CERN-SPSC-2012-031](#)
  - Positive feed-back from SPSC in jan. 2013 : [CERN-SPSC-2013-008](#)
- **Current think tank**
  - F. Arleo, E.G. Ferreira, F. Fleuret, P.-B. Gossiaux, S. Peigné
  - Need people (experimentalist) to pursue investigation
- **apparatus**
  - **Tracking**
    - Needs low detector occupancy → silicon technology
    - **Welcomes group with expertise !**
  - **Calorimetry**
    - Need ultragranular calorimetry à la CALICE
    - Expertise at LLR - Ecole polytechnique (France)
  - **Trigger**
    - Instrumented (magnetized) Fe Absorber
    - **Welcomes group with expertise !**
- **Timeline**
  - From  $T_0$  (3 labs involved): ~ 5 Years for full simulation and final design (2 years), construction and installation (2 years), commissioning (1 year)


WE NEED YOU !!  
JOIN US !



# backup


- **How to Test sequential suppression with charmonia ?**

- **must measure  $J/\Psi$ ,  $\Psi'$ ,  $\chi_c$**
- $\sim 30\%$  (resp.  $\sim 10\%$ ) of inclusive  $J/\Psi$  comes from  $\chi_c$  (resp.  $\Psi'$ ) decay.
- According to lattice calculations,  $T_d(\Psi') < T_d(\chi_c) < T_d(J/\Psi)$
- **If screening, one should observe a step-like suppression patterns**



- **Alternative (no QGP) scenario: suppression by comoving hadrons**

- Smooth suppression
- Same suppression-starting point
- Slopes related to binding energy :  $S_{\Psi'} > S_{\chi} > S_{J/\Psi}$



- Anomalous suppression  
at SPS

[Eur.Phys.J.C49:559-567,2007](#)

## Expectations in comovers scenario

### Binding energy

state	$\eta_c$	$J/\psi$	$\chi_{c0}$	$\chi_{c1}$	$\chi_{c2}$	$\psi'$
mass [GeV]	2.98	3.10	3.42	3.51	3.56	3.69
$\Delta E$ [GeV]	0.75	0.64	0.32	0.22	0.18	0.05


Taking breakup cross-sections:

- comovers-direct  $J/\Psi = 0.2$  mb
- comovers -  $\chi_c = 1.0$  mb
- comovers -  $\Psi' = 2.0$  mb





60% direct  $J/\Psi$   
 + 30%  $\chi_c \rightarrow J/\Psi + \gamma$   
 + 10%  $\Psi' \rightarrow J/\Psi + X$   
**Inclusive  $J/\Psi$  yield**

and considering  
feed-downs



L (fm)	4.37	4.90	6.65	7.65	8.83	9.43
$\epsilon$ (GeV/fm <sup>3</sup> )	1.04	1.24	2.04	2.53	3.19	3.76

# J/Ψ@SPS .vs. Υ@LHC




↑ Charmonia @ SPS

↑ bottomonia @ LHC


Testing sequential suppression scenario at SPS is crucial to fully understand RHIC and LHC results.

# Energy scan

- Spectrometer acceptance: two detector configurations



**Mid-rapidity**  
 $y_{\text{CMS}} \in [-0.5 ; 1]$   
 for  $P_{\text{beam}} = 158 \text{ GeV}/c$



**Forward-rapidity**  
 $y_{\text{CMS}} \in [0.5 ; 2]$   
 for  $P_{\text{beam}} = 158 \text{ GeV}/c$

Depending on the beam energy, different rapidity ranges accessible

$P_{\text{beam}}$ (GeV/c)	$\sqrt{s}$ (GeV)	Rapidity of Center-of-mass	Mid-rapidity		Forward-rapidity	
			$y_{\text{CMS}} \text{ min}$	$y_{\text{CMS}} \text{ max}$	$y_{\text{CMS}} \text{ min}$	$y_{\text{CMS}} \text{ max}$
158	17.2	2.91	-0.5	1	0.5	2
120	15.1	2.77	-0.36	1.14	0.65	2.14
80	12.3	2.57	-0.16	1.34	0.84	2.34
60	10.7	2.43	-0.02	1.48	0.98	2.48



**Common coverage:  $y_{\text{CMS}} \in [0;2]$  (NA50/NA60 coverage =  $[0;1]$ )**

# Rapidity coverage

- A thorough p+A program
  - mandatory as reference for hot nuclear matter effects

$J/\Psi, \Psi', \chi_c$  in a large  $y_{CMS}$  range  
 → Large coverage in  $x_2$   
 → Large coverage in  $x_F$

$E_{beam}$ ( $\sqrt{s}$ )	Exp.	$y_{CMS}$	$x_2$	$x_F$
158 GeV (~17 GeV)	NA50	[0;1]	[0.07;0.18]	[0;0.42]
	CHIC	[-0.5;2]	[0.02;0.30]	[-0.19;1]
450 GeV (~29 GeV)	NA50	[-0.4;0.6]	[0.06;0.16]	[-0.09;0.14]
	CHIC	[-0.9;1.6]	[0.02;0.26]	[-0.22;0.51]



F. Arleo and S. Peigné, arXiv:1204.4609

- CHIC expected performances for low mass dileptons

- Tracking performed upstream to the absorber

- → **no multiple scattering** due to absorber

- → momentum resolution affected by magnetic field only:

$$\frac{\Delta P}{P} \propto \frac{1}{BL^2} P$$

- Momentum resolution

- With a 1m long 2.5T dipolar magnetic field

- $\frac{\Delta P_\mu}{P_\mu} = 1\%$  for typical muon from  $J/\Psi$  ( $\langle P_\mu \rangle \sim 10$  GeV/c)

- $\frac{\Delta P_\mu}{P_\mu} = 0.7\%$  for typical muon from  $\omega$  ( $\langle P_\mu \rangle \sim 7$  GeV/c)

- Expected mass resolution:

- $J/\Psi$ :  $\frac{\Delta P_\mu}{P_\mu} = 1\% \Rightarrow \frac{\Delta P_\mu}{\sqrt{2}P_\mu} = \frac{\Delta M_{\mu\mu}}{M_{\mu\mu}} = 0.7\%$

- $\omega$ :  $\frac{\Delta P_\mu}{P_\mu} = 0.7\% \Rightarrow \frac{\Delta P_\mu}{\sqrt{2}P_\mu} = \frac{\Delta M_{\mu\mu}}{M_{\mu\mu}} = 0.5\%$

$$\Delta M_{\mu\mu}^{J/\Psi} \sim 3.097 \text{ GeV}/c^2 \times 0,7\% \sim 20 \text{ MeV}/c^2$$

$$\text{NA50: } \Delta M_{\mu\mu}^{J/\Psi} \sim 90 \text{ MeV}/c^2$$

$$\Delta M_{\mu\mu}^{\omega} \sim 782.7 \text{ MeV}/c^2 \times 0,5\% \sim 4 \text{ MeV}/c^2$$

$$\text{NA60: } \Delta M_{\mu\mu}^{\omega} \sim 20 \text{ MeV}/c^2$$



## Conclusions of the CERN Town meeting on “Relativistic Heavy-Ion Collisions”



CERN - june 29, 2012

“...The town meeting also observed that the **CERN SPS** would be well-positioned to contribute decisively and at a competitive time scale to central open physics issues at large baryon density. In particular, the CERN SPS will remain also in the future the only machine capable of delivering, heavy ion beams with energies exceeding 30 GeV/nucleon, and **the potential of investigating rare penetrating probes at this machine is attractive.**”

## Results from CMS

### "Observation of Sequential $\Upsilon$ Suppression in PbPb collisions" (at LHC)

*PRL 109, 222301 (2012)*






Charmonia @ SPS     bottomonia @ LHC

**Testing sequential suppression scenario at SPS is crucial to fully understand RHIC and LHC results.**

NA50, PbPb  $\sqrt{s_{NN}}=17.2$  GeV



■ Ψ(1S)  
■ Ψ(2S)



Charmonia @ SPS

bottomonia @ LHC

Testing sequential suppression scenario at SPS is crucial to fully understand RHIC and LHC results.



↑ Charmonia @ SPS

↑ bottomonia @ LHC

**On the same axis**