


Review of recent results
on jet physics in Heavy Ion
from LHC
(compared to RHIC)

Alexandre SHABETAI
CNRS/IN2P3 - SUBATECH (Nantes)

Physics motivation



Jets are produced very early and are sensitive to **early stage** of the collision

→ This allows to **probe and study the QGP** by using **jet properties**

- pp: Study jet **production** (ex. Cross section measurements) : test pQCD
- pA / dA: Are they affected by Cold Nuclear Mater effects ?
- A-A Study in medium **energy loss**
 - **Path length** dependence
 - **Broadening** of shower
 - Leading hadron vs. softening of FF
 - Probe ex. the **density** of the medium

This can be studied by using various observables, in this talk we will mainly discuss: **nuclear modification factors, h-jet azimuthal correlations, di-jet asymmetry and jet FF.**

Nuclear Modification Factors


$$R_{AA} = \frac{\text{Yield}(AA)}{\text{Yield}(pp)}$$

$$R_{AA}(p_T) = \frac{1}{N_{\text{coll}}} \times \frac{dN_{AA}/dp}{dN_{pp}/dp}$$


In absence of nuclear modifications,
hard processes are expected to follow
 N_{col} scaling $\rightarrow R_{AA} = 1$

Single particle RpPb & RdA

RHIC: $R_{dAu} > 1$ at high p_T
 \Rightarrow CNM effects (Cronin / Shadowing)




LHC: High- p_T charged particles follow **binary scaling**.
Initial state effects are small(er)




Tension at high p_T between ALICE and ATLAS/CMS

ALICE / CMS difference
 Likely due to pp reference
CMS: new pp@5 TeV
 interpolation (used for Jet FF)



Issue solved?





single particle R_{AA} at high p_T

arXiv: 1504.04337

60

arXiv:1506.07287

PRC 87 (2013) 034911



ALI-PUB-93398

LHC:

$R_{AA} = 0.55 \pm 0.01(\text{stat}) \pm 0.04(\text{syst})$.


A lot of models describing the trend of R_{AA} with different set of parameters
 R_{AA} not enough to really constrain the transport coefficients

LHC RUN 2: evolution of the shape at higher p_T ?

RHIC:

Similar R_{AA} values in central collisions at LHC and RHIC: with a much harder spectrum at LHC
 → larger ΔE at LHC

Jets Measurements



“Cone like” or sequential reconstruction algorithms (k_T and anti- k_T are used in the following analyses).

Jet production cross-sections in pp collisions




● RHIC : STAR & PHENIX

● ALICE: Full jets

● ATLAS

Phys.Rev. D86 (2012) 014022

PLB 722, 262 2013




Good agreement with NLO pQCD over a broad kinematic range

Important reference for p-Pb & Pb-Pb analyses

Jet R_{pA} @ LHC



$$R_{pA}^{jet} = \frac{dN_{pA}^{jet}/dp_T^{jet}}{N_{coll} dN_{pp}^{jet}/dp_T^{jet}} \cdot \frac{N_{event}^{pp}}{N_{event}^{pA}}$$



jet yield in p-Pb compatible with what is expected
from a **superposition of independent pp collisions**


Consistent with no CNM effects

Results from nucleus nucleus collisions



Background and its fluctuations in Pb-Pb

Ex of ALICE



$$\rho = \text{median} \left(\frac{p_T^{\text{jet},i}}{A_i^{\text{jet}}} \right)$$

→ Background density scales with event multiplicity

$$\rho \sim N \langle p_T \rangle$$

→ $\rho = 200$ GeV/c for most central collisions PbPb collisions at LHC

→ ρ has large event by event up/down fluctuations

Ex: $\sigma_{\text{ch}} \approx 10$ GeV/c for $R=0.4$

→ Smaller $R \rightarrow$ less background fluctuation

→ Limit R to reasonable values $0.2 < R < 0.5$

→ jets studies are challenging.... In heavy ions Especially at low p_T

Larger background fluctuations (σ) for larger R , while larger R should be preferred to recover as much information as possible of the jet property.

Jet Nuclear Modification Factor

$$R_{AA} = \frac{\text{Yield}(AA)}{\text{Yield}(pp)}$$


$$R_{AA}(, p_T) = \frac{1}{N_{\text{coll}}} \times \frac{dN_{AA}/dp}{dN_{pp}/dp}$$

In absence of nuclear modifications,
hard processes are expected to follow
 N_{col} scaling $\rightarrow R_{AA} = 1$

ALICE: Full jet spectrum & R_{AA} in PbPb collisions

- Full jet spectrum in Pb-Pb: large and fluctuating background (JHEP03 (2012) 053)


$$p_{T,jet} = p_{T,rec} - \rho A$$



Phys.Lett. B746 (2015) 1

Strong suppression observed in spectra and R_{AA}

- Full jet R_{AA} :



Phys.Lett. B746 (2015) 1

Leading track p_T cut \rightarrow fragmentation bias

- Suppression quantified by R_{AA} for different centralities:
- 0-10% $R_{AA} \sim 0.28 \pm 0.04$
- 10-30% $R_{AA} \sim 0.35 \pm 0.04$
- Both models use a fit to hadron R_{AA} to adjust their parameters.


Both models are in agreement with data (slight deviation of YaJEM at high p_T in central collisions)

jet RAA: LHC vs RHIC

LHC:

- Full jet R_{AA} in central Pb-Pb collisions,
 - Different jet reconstruction technics (ATLAS: Calo Jets, CMS: PF jets, ALICE: Ch+En Jets) used by the different experiments
 - $R=0.2$
- NB: ATLAS scaled from $R=0.4$ to $R=0.2$

Results at LHC are in fair agreement




RHIC:

Smaller suppression of inclusive jet yield at RHIC


Jets are also less suppressed than hadrons at RHIC

Jet RAA expected to be one if all the jet energy is measured


Where did the energy go ?



PHENIX: Jet RAA



For central collisions, jets are suppressed by about a factor 2



Surface bias effect: the parton producing the jet is biased towards higher in-medium path length
Trigger Hadron: close to the surface

Hadron-jet Azimuthal Correlations

Recoil jet measurement provides us with a good handle on the combinatorial background and allows to go to larger R

Bonus: No fragmentation bias on recoil side


ALICE vs STAR: Hadron-jet correlations

ALICE:

Low $p_{T,\text{trig}}$ (TT*[8, 9]) trigger recoil jet spectrum as a reference (dominated by combinatorial jet).


High $p_{T,\text{trig}}$ (TT[20, 50]) trigger recoil jet spectrum mainly from hard (high Q^2) process (signal).

Δ_{Recoil} is defined as the difference of these two spectra to remove bkg and uncorrelated component. * TT for triggered track



Shape does not change with $p_{T,\text{trig}}$
→ Combinatorial background

$p_{T,\text{jet}} > 20$ GeV/c:
Recoil jet spectrum evolves with $p_{T,\text{trig}}$




STAR: Mixed events describe the combinatorial background
→ Trigger-correlated jet distribution: subtract ME from data
Comparable to ALICE h+jet


Hadron-jet correlations at RHIC

ALICE

arXiv:1506.03984



STAR



STAR: Nuclear modification factor “ I_{CP} ”:

- close to 1 at low p_T
- large suppression at $p_T > 10$ GeV/c: $I_{CP} \sim 0.2$

Larger suppression at RHIC compared to LHC
However: different kinematics and cuts

Constant $\Delta E \sim 8$ GeV




ALICE: Recoil jet yields are suppressed (~ 0.6) (conditional yield) independently of R and slowly decreases with jet p_T

Cone size dependence


First step towards measuring **internal jet structure**
 Measure jets **with different resolution parameters**

ALICE: 1506.03984

ATLAS: Phys. Rev. C 86, 014907 (2012)



No evidence for jet broadening for $R < 0.5$




Excess observed for $R = 0.5$

Jet acoplanarity: Inter jet broadening in p-Pb


Sources of acoplanarity in pp: intrinsic k_T , 3-jet events, hard FSR, ISR

Additional sources in p-Pb: interaction of the partonic projectile with the nuclear medium.

$$k_T = p_{T, \text{ch jet}}^{\text{trigger}} \sin(\Delta\varphi_{\text{dijet}})$$



STAR: small k_T broadning




ALICE : charged jets

No modification of di-jet k_T

medium induced acoplanarity



Large angle inter-jet scattering




D’Eramo, Rajagopal, ‘12
updated calculation underway

ALICE : Width (σ) consistent in Pb+Pb with
PYTHIA embedded data

➔ No evidence of medium-induced
acoplanarity of recoil jets



Au+Au 60-80%






+ Au+Au 0-10%

STAR: High- p_T trigger: no broadening with
central

Low- p_T trigger: broadening with centrality²⁰

➔ Broadening in central collisions at RHIC for
very low p_T jets? Work ongoing

Di-jets Asymmetry



$$A = \frac{(pT_1 - pT_2)}{(pT_1 + pT_2)}$$





Di-jet asymmetry in Pb+Pb collisions

ATLAS: PRL 105 (2010) 252303


First heavy ion jet measurements at LHC

$$A = \frac{(pT_1 - pT_2)}{(pT_1 + pT_2)}$$

di-Jet asymmetry observed in central Pb+Pb collisions by ATLAS and CMS





CMS: PRC 84 (2011)024906



Assymetry : Probing differences in quenching between the two parton showers


ATLAS: x_J

$$x_J = \text{subleading jet } p_T / \text{leading jet } p_T$$



- ATLAS has **fully unfolded** detector resolution effects for dijet p_T asymmetry
- **Difference** between Pb+Pb and pp **get smaller with leading jet p_T** (independent on p_T in CMS)

CMS: Where did the energy go?




courtesy Y. J. Lee (CMS)

Missing p_T


$$p_T^{\parallel} = \sum_i -p_T^i \cos(\phi_i - \phi_{\text{Dijet}})$$

- increases with A_J and is balanced by:
2-8 GeV/c particles in pp
 $p_T < 2$ GeV/c particles in 0-10% Pb+Pb
- high p_T imbalance at small R compensated by low p_T





particles in subleading jet direction extending to **large ΔR**

STAR: di-jets: properties of radiated energy



$$A_J = \frac{p_T^{\text{Lead}} - p_T^{\text{SubLead}}}{p_T^{\text{Lead}} + p_T^{\text{SubLead}}}$$



Find jet with high p_T cut on constituents → reduce combinatorial background
rerun jet finder for identified jets with looser p_T cut or different radius

Momentum balance restored with $R=0.4$, $p_T^{\text{cut}} = 0.2$ GeV

Imbalance with:






- increasing p_T^{cut} → Jet softening
- decreasing radius → jet broadening (between 0.2 and 0.4)

Different modification of jet structure at RHIC and LHC

BUT Non negligible jet p_T biases possible (cuts & low p_T jets → background?)



ALICE : Jet v2



Qualitative **agreement** between ALICE and ATLAS

Good agreement with **JEWEL** in **peripheral** collisions

Indication of non-zero jet v2 as a result of fluctuations in initial participant density

→ large parton energy loss, sensitive to collision geometry up to high pT

ALICE: arXiv:1509.07334
ATLAS: PRL 111 152301 (2013)



ATLAS: Path length effects on di-jet production



Goal : constrain path length dependence of energy loss

ATLAS-CONF-2015-021

Previous study of jet production vs. event plane showed **modest path length dependence** ($v_{2, \text{jet}} \sim 2-5\%$)

ATLAS: PRL 111 (2013)152301


$$\langle A_J \rangle = A_J^0 \left(1 + 2c_2^{\text{obs}} \cos(2(\phi^{\text{lead}} - \Psi_2)) \right)$$



→ c_2 is small (<2%) and negative
 → larger A_J for leading jets oriented out-of-plane than for in-plane ones

Study un-equal path lengths of the showers in the medium

- Selecting on the **angle between the leading jet and 2nd order Event-Plane angle Ψ_2** , provides control on the path length traversed by the jet pair.
- Jet pairs that are **Out-of-Plane** are expected to **traverse more medium** than In-Plane.




Extract c_2 that quantifies the EP angle dependence of A_J



ATLAS: Neighbouring jet production in Pb-Pb collisions

Annulus around the test jet

Neighbouring jet



Goal: probe differences in quenching that do not primarily result from difference in path length. (fluctuations of energy loss)

Production of neighbouring jets quantified using a rate of jets accompanying a given test jet

$$R_{\Delta R} = \frac{1}{dN_{\text{jet}}^{\text{test}}/dE_T^{\text{test}}} \sum_{i=1}^{N_{\text{jet}}^{\text{test}}} \frac{dN_{\text{jet},i}^{\text{nbr}}}{dE_T^{\text{test}}} (E_T^{\text{test}}, E_{T,\text{min}}^{\text{nbr}}, \Delta R)$$

$$\rho_{R \Delta R} = R_{\Delta R}(\text{central})/R_{\Delta R}(\text{peripheral})$$

Study its dependence on:


- E_T of test jet:**

Suppression of neighbouring jet production in central Pb-Pb is:

~ independent of test jet E_T & similar to inclusive jets

- E_T of neighbouring jet**

Nuclear modification factor for neighbouring jets approaches 1 when $E_{T,\text{test}} \approx E_{T,\text{nbr}}$



Fragmentation Functions

Study the momentum distribution of tracks in jets

arXiv:1411.4969

● Charged jets in pp collisions @ 7 TeV:

For $\xi < 2$ **scaling**


At higher ξ , maximum : **'hump-backed plateau'** → **suppression of low momentum particle** production by QCD coherence (angular ordering)

With **increasing jet pT**, the **area** of the distributions **increases** (**higher particle multiplicity** in jets),

maximum shifts to higher values of ξ

This observation is in qualitative agreement with **MLLA**


Measurement challenging in Pb-Pb collisions (large heavy ion background and fluctuations).



ALI-PUB-92483

CMS/ATLAS: FF in pA

Reduce background → high p_T jets needed




CMS: no modification of fragmentation functions (within uncertainties). Coherent with R_{pPb} (jets)=1 if R_{pPb} (charged) will move to ~ 1

ATLAS: suggestion for a hardening of the FF

→ Also comes from the interpolation method used for the pp reference ?

FF in PbPb

Reduce background → high p_T jets needed



$$Z = p_{||\text{Trk}} / p_{\text{Jet}}$$

Qualitative agreement between ATLAS and CMS

Small modifications:



Depletion from 3-4 GeV to 40-50 GeV (2-3% of the total jet energy)

Enhancement below 3-4 GeV ($\sim 2\%$ of the jet energy)

→ Need to improve systematics

CMS: jet shapes (radial profile)

$$\rho(r) = \frac{1}{f_{\text{ch}}} \frac{1}{\delta r} \frac{1}{N_{\text{jet}}} \sum_{\text{jets}} \frac{p_{\text{T}}(r - \delta r/2, r + \delta r/2)}{p_{\text{T}}^{\text{jet}}}$$





- Low- p_{T} excess of jet tracks in sub-leading jet side extended to large angles

Exclusive measurements

Strange particle production in jets

$p_{T, \text{jet}} > 10 \text{ GeV}/c$

$p_{T, \text{jet}} > 20 \text{ GeV}/c$





Λ/K^0 ratio in charged jets in p-Pb and Pb-Pb are significantly lower than for inclusive particles

→ Jet composition is not influenced by the enhanced baryon/meson production in the bulk.

B-jets

PRL 113 (2014) 132301



b-jet R_{AA} decreases with centrality down to ~ 0.4 in 0-10%

Similar suppression to inclusive jets for $p_T > 80$ GeV/c

➔ Mass effects might appear at lower p_T

Summary

A lot of results on jets from both LHC RUN 1 and RHIC

→ In general good consistency between experiments

More **differential observables** are needed in order to better understand and constrain energy loss mechanisms

Theory / Experiments comparisons are very important

→ Are we really comparing the same observables ?

→ Interactions between theorists / experimentalists are crucial.

Outlook

The LHC **run 2** is just starting... (STAR / ALICE **upgrades**)

- Increased luminosity → **more statistics will be available**
 - Redo Run 1 measurements to **reduce the systematics incertenties**
 - **new observables** are under study (ex PID in jets, HF jets, quark / gluon jets tagging, sub-jets, ...)
- many new exiting results **are expected soon** → Stay tuned !

Backup